

THREE NEWLY RECORDED GENERA AND SPECIES OF GOBIID FISHES (TELEOSTEI: GOBIIDAE) FROM THE DONGSHA ATOLL (PRATAS ISLANDS), SOUTH CHINA SEA

Shao-Liang Hsu^{1,2}, Ta-Kang Liu¹, Kuan-Te Chen³, Kwang-Tsao Shao⁴ and I-Shiung Chen³

Key words: Gobiidae, new records, fish taxonomy, fish fauna, Dongsha Atoll, Taiwan.

ABSTRACT

Three newly recorded gobiid genera of coral-reef gobies have been collected and examined by scuba diving in the Dongsha Atoll, also known as the Pratas Islands, South China Sea. They are *Feia* Smith, 1959; *Sueviota* Winterbottom and Hoese, 1988; and *Trimmatom* Winterbottom and Emery, 1981. These three genera were formerly unrecorded in the Dongsha Atoll and all Taiwanese waters before this paper. One representing species was found for each gobiid genus. The diagnosis for these three newly recorded species and genera will be provided in this paper.

I. INTRODUCTION

The generally small body-sized gobiid fishes in Family Gobiidae comprise the most diverse group among marine teleost fishes (Miller [7]). Among them, the tropical and subtropical coral-reef associated gobies are the most diverse group of the species component of gobioid fishes.

In order to investigate the species diversity of gobioid fishes in the South China Sea, our research team has surveyed the fish fauna during the 2011 marine biodiversity expedition to the Dongsha Atoll, also known as the Pratas Islands, especially in the northern and eastern outside slopes of the great lagoon of Dongsha Atoll.

More recently, Chen et al. [4] have published a colored fish ecology guide-book of the Dongsha Atoll, including underwater visual census data of fishes by scuba diving. In their fish checklist there were 24 genera and 74 species of gobiid fishes, but some of them still lack specimen evidence. In addition,

we found three newly recorded genera and species of gobioid fishes that have never been reported in the Atoll and all Taiwanese waters.

The aim of this paper is to document these three newly recorded genera and species of gobiid fishes from the Dongsha Atoll. They are *Feia* Smith, 1959 [13]; *Sueviota* Winterbottom and Hoese, 1988 [25]; and *Trimmatom* Winterbottom and Emery, 1981 [24]. Fish photos are also provided in this paper.

II. MATERIALS AND METHODS

All specimens of the newly recorded species were collected by hand-net during scuba diving. All counts and measurements were made from specimens stored in 70% ethanol after 10% formalin preservation. Morphometric methods follow Miller [7] and meristic methods follow Akihito et al. [1], Chen and Shao [3], Chen et al. [2], and Suzuki et al. [15]. Terminology of cephalic sensory canals and free neuromast organs (sensory papillae) follow Wongrat and Miller [26], based on Sanzo [11]. All fish specimens are deposited at the Pisces collection of National Taiwan Ocean University (NTOUP), Keelung.

Abbreviations: A, anal fin; C, caudal fin; D1 and D2, 1st and 2nd dorsal fins respectively; LR, longitudinal scale rows; P, pectoral fin; PreD, predorsal scales; TR, transverse scale rows; V, pelvic fin. All fish lengths are standard length (SL).

III. TAXONOMY

Family GOBIIDAE

Feia Smith, 1959

Type species: *Feia nympha* Smith, 1959: 206 [13]

Diagnostic characters of genus: (1) five branchiostegal rays; (2) fin rays: D1 VI, D2 I/7-10, AI/7-9, P 14-18; (3) squamation: LR 18-27 (scales only on caudal peduncle varied to covered all lateral trunk); (4) gill-opening restricted, extending only about vertical line at posterior 2/3 region of opercle; and (5) head lateral-line system: no canal pore to reduced canal pattern; typically reduced, longitudinal papillae pattern.

Distribution: Indo-West Pacific.

Remarks: There are 4 nominal, valid species in the Indo-West

Paper submitted 05/11/12; revised 08/27/12; accepted 11/07/12. Authors for correspondence: Kwang-Tsao Shao (e-mail: zoskt@gate.sinica.edu.tw) and I-Shiung Chen (e-mail: isc@ntou.edu.tw).

¹Institute of Ocean Technology and Marine Affairs, National Cheng-Kung University, Tainan 701, Taiwan, ROC.

²Marine National Park Headquarters, Taiwan, Kaohsiung 811, Taiwan, ROC.

³Institute of Marine Biology & CMBB, National Taiwan Ocean University, Keelung 202, Taiwan, ROC.

⁴Biodiversity Research Center, Academia Sinica, Taipei 115, Taiwan, ROC.

Pacific including: the type species *Feia nympha* Smith, 1959 [13] as well as 3 other species: *Feia nota* Gill & Mooi, 1999 [5]; *Feia ranta* Winterbottom, 2003 [22]; and *Feia dabra* Winterbottom, 2005 [23]. The species of genus can exist from 2 m to 25 m water depth of ecological habitats of coral-reef debris, holes and sandy bottom [5, 13, 22, 23].

***Feia nympha* Smith, 1959**

(Fig. 1)

Feia nympha Smith, 1959: 206 [13] (Pinda, Mozambique).
Feia nympha, Gill and Mooi, 1999: 369 [5]; Akihito et al. in Nakabo, 2002: 1249 [8]; Winterbottom, 2003: 101 [22].

Material examined:

NTOUP-2011-06-108, 14.3 mm SL, coll. K.T. Chen et al., June 20, 2011, GPS 20° 43' 52.93" N 116°42' 24.01" E, 15-20 m depth, northwestern region of lagoon, Dongsha Atoll (Pratas Islands), South China Sea, Taiwan.



Fig. 1. *Feia nympha*, NTOUP-2011-06-108, 14.3 mm SL, Dongsha Atoll (Pratas Islands), South China Sea. Photograph by K.T. Chen.

Diagnosis. This species can be well distinguished from other congeners by the following unique combination of features:

- (1) **Fin rays:** D1 VI; D2 I/9; A I/8; P 14; V I/5. D1 without filamentous rays.
- (2) **Squamation:** LR 20; PreD 0. Scales extending anteriorly to middle of body. Head, predorsal region and anterior part of trunk naked.
- (3) **Gill-opening:** restricted, extending only to vertical line at posterior 2/3 region of opercle.
- (4) **Body proportion:** all morphometric data shown in Table 1.
- (5) **Head lateral-line system: Head canals:** No head canal pores. **Sensory papillae:** Overall infraorbital papillae pattern. Rows *b* and *d* with densely set papillae. Row *f* as long longitudinal papilla row.
- (6) **Colouration:** Body generally creamy yellow, with many tiny irregular brown spots. Dorsum with several brown cross-bars. Caudal fin base with a larger brown spot. Dorsal half of head with tiny rows of brown spots, ventral half of head yellowish. Second dorsal fin with many oblique brown spots. Pectoral fin base yellowish. Caudal fin translucent with many tiny brown spots.

Distribution:

Indo-West Pacific, including the Red Sea. This species was first

found in Dongsha Atoll (Pratas Islands), South China Sea. However, it has not yet been found in Taiwanese waters.

***Sueviota lachneri* Winterbottom and Hoese, 1988**

Type species: *Sueviota lachneri* Winterbottom and Hoese, 1988 [25]

Diagnostic characters of genus: (1) five branchiostegal rays; (2) fin rays: D1 VI, D2 I/8-10, A I/7-9, P 16-19; (3) squamation: LR 24-27 (scales covered all lateral trunk), predorsal region entirely naked; (4) gill-opening moderate, extending ventrally to about vertical line at midline of opercle; and (5) head lateral-line system: no canal pore to reduced canal pattern with presence of both anterior and posterior oculoscapular canals, but lacking preopercular canal; typically brief and reduced, longitudinal papillae pattern.

Distribution: Indo-West Pacific.

Remarks: There are 4 nominal, valid species in the Indo-West Pacific including: the type species: *Sueviota lachneri* Winterbottom & Hoese, 1988 [25] as well as 3 other species: *Sueviota aprica* Winterbottom & Hoese, 1988 [25], *Sueviota atrinasa* Winterbottom & Hoese, 1988 [25] as well as *Sueviota larsonae* Winterbottom & Hoese, 1988 [25]. The most species of genus can exist from 8 m to 48 m water depth of ecological habitats of coral-reef caves and holes except the deeper species, *S. larsonae* found from 40 m up to 82 m depth [25]. Sunobe and Shinomyia [14] have ever documented the reproduction behavior of *S. lachneri* as “the entire reproductive sequence consisted of nest preparation by a male, courtship behaviors, nest entry by a female, spawning, and paternal care until the hatching of eggs”.

***Sueviota lachneri* Winterbottom and Hoese, 1988**

(Fig. 2)

Sueviota lachneri Winterbottom and Hoese, 1988: 9 [25] (Drop-off on east side of Isle Poule, Salomon Atoll, Chagos Archipelago, central Indian Ocean).

Sueviota lachneri, Nakabo, 2000: 1184 [8]; Akihito et al. in Nakabo, 2002: 1184 [8]; Randall, 2005: 552 [9]; Hoese and Larson, 2006: 1685 [6].

Material examined:

NTOUP-2011-07-222, 3 specimens, 13.6-16.2 mm SL, coll. I-S. Chen et al., July 7, 2011, GPS 20°38' 30.21" N 116°54' 47.42" E, 10-15 m depth, eastern outside slope of great lagoon, Dongsha Atoll (Pratas Islands), South China Sea, Taiwan.



Fig. 2. *Sueviota lachneri*, NTOUP-2011-07-222, 16.2 mm SL, Dongsha Atoll (Pratas Islands), South China Sea. Photograph by K.T. Chen.

Diagnosis.

This species can be well distinguished from other congeners by the following unique combination of features:

(1) **Fin rays:** D1 VI; D2 I/9; A I/8; P 16; V I/5. Rear tips of D2 and A do not extend to procurent rays of C.

(2) **Squamation:** LR 25–26; TR 6; PreD 0.

(3) **Gill-opening:** Moderate, extending ventrally to vertical line at midline of opercle.

(4) **Body proportion:** All morphometric data shown in Table 1.

(5) **Head lateral-line system: Head canals:** Anterior oculo-scapular canal present with singular pore λ and κ , terminating at pore α but lacking pore β . Posterior oculoscapular canal with two terminal pores γ and ϵ . No preopercular canal. **Sensory papillae:** Rather brief infraorbital longitudinal papilla pattern. Both longitudinal papilla rows *a* and *b* with 2–4 papillae.

(6) **Colouration:** Body generally pale to light pinkish. Anterior nostrils orange red with distal black rim. Iris and both lips orange red. Cheek with several transverse orange red bars. Pectoral fin with two horizontal orange red bars. Seven inner transverse orange red bands from anal fin origin to caudal fin base. Dorsals with several rows of orange red spots. Caudal fin translucent with 4 main vertical rows of red spots. Anal fin grayish.

Distribution:

Indo-West Pacific. This species was first found in Dongsha Atoll (Pratas Islands), South China Sea.

***Trimmatom* Winterbottom and Emery, 1981**

Type species: *Trimmatom nanus* Winterbottom and Emery, 1981 [24]

Diagnostic characters of genus: (1) five branchiostegal rays; (2) fin rays: D1 VI, D2 I/8–12, A I/7–11, P 15–19; (3) LR 0–27 (body no scales varied to body scales covered all lateral trunk), predorsal region entirely naked; (4) gill-opening rather wide, extending ventrally forward about vertical line slightly beyond preopercle; and (5) head lateral-line system: no any head canal and pore; typically brief and reduced, longitudinal papillae pattern.

Distribution: Indo-West Pacific.

Remarks: There are 7 nominal, valid species in the Indo-West Pacific including: the type species *Trimmatom nanus* Winterbottom and Emery, 1981 [24] as well as 6 other species: *Trimmatom macropodus* Winterbottom, 1989 [16], *Trimmatom eviotops* (Schultz, 1943) [12], *Trimmatom offucius* Winterbottom & Emery, 1981 [24], *Trimmatom pharus* Winterbottom, 2001 [20], *Trimmatom sagma* Winterbottom, 1989 [16], and *Trimmatom zapotes* Winterbottom, 1989 [16]. The species member of genus (the type species) comprises a smallest coral-reef benthic fish in the World [24]. The species of genus

mostly exist from 3 m to 50 m water depth of ecological habitats of coral-reef caves and holes [16, 20, 24].

***Trimmatom nanus* Winterbottom and Emery, 1981** (Fig. 3)

Trimmatom nanus Winterbottom and Emery, 1981: 143 [24] (Salomon Atoll, Chagos Archipelago, northern Indian Ocean, 5°18'04" S, 72°15'44" E, depth 18–25 meters). *Trimmatom nanus*, Winterbottom, 1989: 1405 [16]; Winterbottom, 1990: 254 [17]; Winterbottom, 2001: 20 [18]; Winterbottom, 2002: 49 [21]; Randall et al., 2002: 162 [10]; Randall, 2005: 555 [9]; Hoese and Larson, 2006: 1691 [6].

Material examined:

NTOUP-2011-07-203, 4 specimens, 7.1–8.7 mm SL, coll. I-S. Chen et al., July 7, 2011, GPS 20°38' 30.21" N 116°54' 47.42" E, 12–15 m depth, eastern outside slope of great lagoon, Dongsha Atoll (Pratas Islands), South China Sea, Taiwan.



Fig. 3. *Trimmatom nanus*, NTOUP-2011-07-203, 8.7 mm SL, Dongsha Atoll (Pratas Islands), South China Sea. Photograph by K.T. Chen.

Diagnosis.

This species can be well distinguished from other congeners by the following combination of features:

(1) **Fin rays:** D1 VI; D2 I/9; A I/9; P 16; V I/5.

(2) **Squamation:** LR 0; PreD 0. Entire head and body naked.

(3) **Gill-opening:** Rather wide, extending ventrally forward to vertical line slightly beyond preopercle.

(4) **Body proportion:** All morphometric data shown in Table 1.

(5) **Head lateral-line system: Head canals:** No head canal pores. **Sensory papillae:** Rather brief infraorbital longitudinal papilla pattern. Both longitudinal papillae rows *a* and *b* with 2–3 papillae. Row *p* surrounding orbit, toward dorsal side of snout.

(6) **Colouration:** Body with dense black melanophores on anterior body. Generally red in adults but purple black in young individuals. Some translucent marks arranged dorsally and ventrally. Iris bright red. Second dorsal, anal and anal fins translucent with red unbranched rays in adults.

Distribution:

Indo-West Pacific, including the Red Sea. This species was first found in Dongsha Atoll, South China Sea. However, it has not yet been found in Taiwanese waters.

Remarks:

This species is still one of the smallest, benthic, coral-reef associated fishes in the world (Winterbottom and Emery [24]). Our specimens collected here provide the northernmost record of this species.

Table 1. Morphometry of three newly recorded gobiid fishes in Dongsha Atoll, South China Sea

<i>Feia nympha</i>	
Cat. No.	NTOUP-2011-06-108
No. of specimen(s)	1
Standard length (SL)	14.3 mm
% in SL	
Head length	30.4
Predorsal length	42.0
Snout to 2nd dorsal length	62.4
Snout to anus	62.1
Snout to anal fin origin	68.8
Prepelvic length	30.2
Caudal peduncle length	13.8
Caudal peduncle depth	13.4
1st dorsal fin base	16.0
2nd dorsal fin base	24.6
Anal fin base	20.4
Caudal fin length	37.4
Pectoral fin length	28.8
Pelvic fin length	20.7
Body depth at pelvic fin origin	14.0
Body depth at anal fin origin	15.7
Body width at anal fin origin	10.7
Pelvic fin origin to anus	31.4
% in HL	
Snout length	32.0
Eye diameter	17.5
Cheek depth	31.1
Postorbital length	63.4
Head width at maximum	69.1
Head width at upper gill-opening	45.9
Bony interorbital width	10.6
Fleshy interorbital width	23.3

Lower jaw length	43.5
------------------	------

Sueviota lachneri

Cat. No.	NTOUP-2011-07-222
No. of specimen(s)	3
Standard length (SL)	13.6 – 16.2 mm
% in SL	
Head length	30.3 – 31.2(30.8)
Predorsal length	37.7 – 38.3(38.0)
Snout to 2nd dorsal length	59.4 – 61.2(60.1)
Snout to anus	57.0 – 60.7(58.8)
Snout to anal fin origin	63.9 – 65.7(65.0)
Prepelvic length	30.6 – 32.1(31.5)
Caudal peduncle length	16.5 – 20.9(18.9)
Caudal peduncle depth	13.0 – 13.3(13.2)
1st dorsal fin base	17.1 – 19.1(18.0)
2nd dorsal fin base	22.7 – 26.0(24.5)
Anal fin base	18.3 – 20.1(19.2)
Caudal fin length	32.6 – 35.1(33.8)
Pectoral fin length	29.1 – 36.2(32.8)
Pelvic fin length	32.1 – 35.6(34.4)
Body depth at pelvic fin origin	19.7 – 21.8(20.8)
Body depth at anal fin origin	18.1 – 19.9(18.9)
Body width at anal fin origin	8.7 – 9.5(9.1)
Pelvic fin origin to anus	28.3 – 31.6(29.9)
% in HL	
Snout length	25.1 – 25.7(25.4)
Eye diameter	28.0 – 30.8(29.8)
Cheek depth	32.2 – 35.1(33.3)
Postorbital length	61.2 – 64.3(62.4)
Head width at maximum	57.5 – 60.2(59.3)
Head width at upper gill-opening	34.7 – 38.8(36.3)
Bony interorbital width	2.3 – 3.1(2.8)
Fleshy interorbital width	11.7 – 14.5(13.3)
Lower jaw length	44.0 – 48.2(46.2)

Trimmatom nanus

Cat. No.	NTOUP-2011-07-203
No. of specimen(s)	4
Standard length (SL)	7.1 – 8.7 mm

% in SL	
Head length	27.7 – 29.9(28.7)
Predorsal length	42.5 – 44.1(43.6)
Snout to 2nd dorsal length	61.4 – 63.1(62.3)
Snout to anus	57.7 – 61.1(59.2)
Snout to anal fin origin	61.2 – 64.4(62.7)
Prepelvic length	28.5 – 32.4(30.2)
Caudal peduncle length	13.4 – 16.0(14.6)
Caudal peduncle depth	11.3 – 12.4(11.8)
1st dorsal fin base	17.6 – 19.2(18.7)
2nd dorsal fin base	24.7 – 30.0(27.5)
Anal fin base	18.8 – 23.5(22.1)
Caudal fin length	27.8 – 30.7(28.7)
Pectoral fin length	21.1 – 28.2(25.0)
Pelvic fin length	36.7 – 39.9(37.7)
Body depth at pelvic fin origin	17.6 – 23.6(19.2)
Body depth at anal fin origin	14.3 – 17.9(15.7)
Body width at anal fin origin	7.3 – 9.0(8.1)
Pelvic fin origin to anus	28.0 – 31.5(30.1)
% in HL	
Snout length	20.8 – 22.8(22.2)
Eye diameter	35.7 – 39.1(37.0)
Cheek depth	31.5 – 35.0(32.8)
Postorbital length	54.9 – 61.9(57.6)
Head width at maximum	50.6 – 58.4(53.7)
Fleshy interorbital width	8.0 – 11.8(9.6)
Lower jaw length	42.1 – 49.1(46.4)

ACKNOWLEDGEMENTS

The authors wish to thank the government research grant support from the National Science Council, Taiwan and also Marine National Park Headquarters, which is responsible for the management of Dongsha Atoll National Park. We are also very grateful to the Center of Excellence for Marine Bioenvironment and Biotechnology, National Taiwan Ocean University for the kind partial grant support for gobioid research in the South China Sea.

REFERENCES

- Akihito, Prince., Hayashi, M., Yoshino, T., Shimada, K., Senou, H. and Yamamoto, T., "Suborder Gobioidi." in Masuda, H., Amaoka, K., Araga, C., Uyeno, T., Yoshino, T. editors. *The fishes of Japanese Archipelago, English text and plates*. Tokai University Press, Tokyo, pp. 228-289 (1984).
- Chen, I-S., Suzuki, T., Senou, H., "A new species of gobioid fish, *Luciogobius* from Ryukyus, Japan (Teleostei: Gobiidae)." *Journal of Marine Science and Technology, Taiwan*, Vol. 16, pp. 250-254 (2008).
- Chen, I-S. and Shao K.T., "A taxonomic review of the gobioid fish genus *Rhinogobius* Gill, 1859, from Taiwan, with descriptions of three new species." *Zoological Studies*, Vol. 35, pp. 200-214 (1996).
- Chen, J.P., Jan, R.Q., Hwang, J.H., Kou, R.W. and Shao, K.T., Fishes of Dongsha Atoll in South China Sea. Marine National Park, Kaohsiung (2011). (in Chinese)
- Gill, A.C. and Mooi, R.D., "*Feia nota*, a new species of gobioid fish from Western Australia." *Records of the Western Australian Museum*, Vol. 19, pp. 365-370 (1999).
- Hoesel, D.F. and Larson, H.K., "Gobiidae." In: *Zoological Catalogue of Australia. Volume 35. Fishes*, pp. 1612-1697 (2006).
- Miller, P.J., "New species of *Corcyrogobius*, *Thorogobius*, and *Wheelerigobius* from West Africa (Teleostei: Gobiidae)." *Journal of Natural History*, Vol. 22, pp. 1245-1262 (1988).
- Nakabo, T., (Ed.) *Fishes of Japan with pictorial keys to the species, English edition*. Tokai University Press, Tokyo, Second edition, Vol. 2, pp. 867-1749 (2002).
- Randall, J.E., *Reef and shore fishes of the South Pacific. New Caledonia to Tahiti and the Pitcairn Islands*. University of Hawaii Press, Honolulu, pp. 1-707 (2005).
- Randall, J.E., Bacchet, P., Winterbottom, R. and Wrobel, L., "Fifty new records of shore fishes from the Society Islands and Tuamotu Archipelago." *aqua, Journal of Ichthyology and Aquatic Biology*, Vol. 5, pp. 153-166 (2002).
- Sanzo, L., "Distribuzione delle papille cutanee (organi ciatiform) e suo valore sistematico nei Gobi." *Mitteilungen aus der Zoologischen Station zu Neapel*, Vol. 20, pp. 251-238 (1911) (in Italian).
- Schultz, L.P., "Fishes of the Phoenix and Samoan islands collected in 1939 during the expedition of the U. S. S. "Bushnell." *Bulletin of the United States National Museum*, Vol. 180, pp. 1-316 (1943).
- Smith, J.L.B., "Gobioid fishes of the families Gobiidae, Periophthalmidae, Trypauchenidae, Taenioididae and Kraemeriidae of the western Indian Ocean." *Ichthyological Bulletin of the J. L. B. Smith Institute of Ichthyology*, Vol. 13, pp. 185-225 (1959).
- Sunobe, T. and Shinomiya, A., "Reproductive behavior, embryonic development and yolk-sac larva of *Sueviota lachneri* (Pisces: Gobiidae) in the laboratory." *Natural History Research (Chiba)*, Vol. 7, pp. 95-99 (2002).
- Suzuki, T., Chen, I-S. and Senou, H., "A new species of *Rhinogobius* Gill, 1859 (Teleostei: Gobiidae) from the Bonin island, Japan." *Journal of Marine Science and Technology, Taiwan*, Vol. 19, pp. 693-701 (2011).
- Winterbottom, R., "A revision of the *Trimmatom nanus* species complex (Pisces, Gobiidae), with descriptions of three new species and redefinition of *Trimmatom*." *Canadian Journal of Zoology*, Vol. 67, pp. 2403-2410 (1989).
- Winterbottom, R., "The *Trimmatom nanus* species complex (Actinopterygii, Gobiidae): phylogeny and progenetic heterochrony." *Systematic Zoology*, Vol. 39, pp. 253-265 (1990).
- Winterbottom, R., "Two new gobioid fish species in *Trimma* and *Trimmatom* (Teleostei: Gobiidae) from the Indian and Western Pacific Oceans." *aqua, Journal of Ichthyology and Aquatic Biology*, Vol. 5, pp. 19-24 (2001).
- Winterbottom, R., "A revision of the *Trimmatom nanus* species complex (Pisces, Gobiidae), with descriptions of three new species and redefinition of *Trimmatom*." *Canadian Journal of Zoology*, Vol. 67, pp. 2403-2410 (1989).
- Winterbottom, R., "Two new gobioid fish species in *Trimma* and *Trimmatom* (Teleostei: Gobiidae) from the Indian and Western Pacific Oceans." *aqua, Journal of Ichthyology and Aquatic Biology*, Vol. 5, pp. 19-24(2001).

21. Winterbottom, R., "Two new species of *Trimma* (Gobiidae) from the central, western, and south Pacific." *aqua, Journal of Ichthyology and Aquatic Biology*, Vol. 5, pp. 45-52 (2002).
22. Winterbottom, R., "*Feia ranta*, a new species of gobiid fish (Acanthopterygii; Perciformes) from Vietnam." *aqua, Journal of Ichthyology and Aquatic Biology*, Vol. 7, pp. 97-102 (2003).
23. Winterbottom, R., "*Feia dabra*, a new species of gobiid fish (Acanthopterygii; Perciformes) from Palau." *aqua, Journal of Ichthyology and Aquatic Biology*, Vol. 10, 45-50 (2005).
24. Winterbottom, R. and Emery, A.R., "A new genus and two new species of gobiid fishes (Perciformes) from the Chagos Archipelago, central Indian Ocean." *Environmental Biology of Fishes*, Vol. 6, pp. 139-149 (1981).
25. Winterbottom, R. and D. F. Hoese., "A new genus and four new species of fishes from the Indo-West Pacific (Pisces; Perciformes; Gobiidae), with comments on relationships." *Royal Ontario Museum Life Sciences Occasional Paper*, Vol. 37, pp. 1-17 (1988).
26. Wongrat, P. and Miller, P.J., "The innervation of head neuromast rows in eleotridine gobies (Teleostei: Gobioidi)." *Journal of Zoology*, Vol. 225, pp. 27-42 (1991).